

*Metretopodidae LESTAGE, 1938

**Metretopus* EATON, 1901

**Metretopus borealis* (EATON, 1871) (LF 14 tr.)

Heptagenia borealis EATON, 1871 (orig.)

Rhithrogena borealis; (comb.) Eaton (1885)

Metretopus norvegicus Eaton, 1901; (typ. gen.) (syn.) Brekke (1940); vgl. Kimmins (1960)

Baetidae LEACH, 1815

Baetis LEACH, 1815

sg. *Baetis* s. str.

Baetis (Baetis) alpinus (PICTET, 1843)

Cloe alpina Pictet, 1843; (orig.)

Cloeon alpina; (comb.) Walker (1853)

Baetis amnicus Eaton, 1871; (syn.) Eaton (1885)

Baetis carpathica Morton, 1910; (syn.) Müller-Liebenau (1969)

Baetis Dorigeri Degrange, 1957a; (syn.) Müller-Liebenau (1969)

Baetis (Baetis) buceratus EATON, 1870

Baetis buceratus Eaton, 1870; (orig.)

Baetis scanicus Bengtsson, 1917; (syn.) Müller-Liebenau (1964)

Baetis Grandii Grandi, 1948b; (syn.) Müller-Liebenau (1969)

Baetis (Baetis) fuscatus (LINNAEUS, 1761)

Ephemera fuscata Linnaeus, 1761; (orig.) (typ. gen.)

Baetis fuscata; (comb.) Stephens (1836)

Cloeon fuscata; (comb.) Walker (1853)

Baetis fuscatus; (comb.) Eaton (1870)

Ephemera bioculata Linnaeus, 1758; (syn.) Kimmins (1964), Brinck & Müller-Liebenau (1965)

Ephemera binoculata auct. = häufige Fehlschreibung von *bioculata*

Ephemera flava von Paula Schrank, 1776; (syn.) Eaton (1871: via *binoculata*)

Ephemera diaphana O.F. Müller, 1776; (syn.) Eaton (1871: via *binoculata*)

? *Ephemera lutea* Fourcroy, 1785 (nomen praeoccupatum) nec Linnaeus; (syn.) Olivier (1791) (via *bioculata*)

Ephemera notata Gmélin, 1789; (syn.) Eaton (1871: via *binoculatus*)

Baetis flavescens Curtis, 1834; (syn. ?) Eaton (1871: via *binoculatus*)

Baetis autumnalis Curtis, 1834; (syn. ?) Eaton (1871: via *binoculatus*)

Baetis venustus Eaton, 1885; (syn.) Müller-Liebenau (1967)

Baetis andalusicus Navás, 1911; (syn.) Alba-Tercedor (1984)

Baetis (Baetis) liebenauae KEFFERMÜLLER, 1974

Baetis liebenauae Keffermüller, 1974 (orig.)

Baetis (Baetis) lutheri MÜLLER-LIEBENAU, 1967

Baetis lutheri Müller-Liebenau, 1967; (orig.)

~ *Baetis subalpinus*: Müller-Liebenau (1960) nec Bengtsson; t. Müller-Liebenau (1967)

Baetis (Baetis) melanonyx (PICTET, 1843)

Cloe melanonyx Pictet, 1843; (orig.)

Cloeon melanonyx; (comb.) Walker (1853)

Baetis melanonyx; (comb.) Eaton (1871)

Baetis kulindrophtalmus Bogoescu, 1933; (syn.) Müller-Liebenau (1969)

Baetis principii Grandi, 1949; (syn.) Müller-Liebenau (1969)

Baetis bengtssoni Müller-Liebenau, 1966; (syn.) Müller-Liebenau (1969)

[= *Baetis finitimus* Eaton, 1871; (syn.) Eaton (1885), falsus = vernus; t. Müller-Liebenau (1969)]

Baetis (Baetis) nexus NAVÁS, 1918

Baetis pentapleobodes Ujhelyi, 1966; (syn.) Alba-Tercedor (2002)

Baetis (Baetis) nubecularis EATON, 1898

Baetis nubecularis Eaton, 1898; (orig.)

**Baetis (Baetis) pavidus* GRANDI, 1951 in Grandi (1951b) (LF 13)

Baetis parva Grandi, 1949 (orig.) (nomen praeoccupatum) [nec *Baetis parvus* Dodds, 1923]

Baetis pavidus; (nomen novum) Grandi (1951b)

Baetis pavidus Grandi, 1949 sensu Grandi, 1953 et auct.